

STABILITY THRESHOLD FOR MULTIADDITIVE AND SYMMETRIC MAPPINGS

Dan M. DĂIANU

Abstract. Z. Gajda showed that the control functions of the form $\epsilon (\|x\|^r + \|y\|^r)$ do not provide stability for additive transformations if and only if $r = 1$. In this note we prove a similar result for n -additive and symmetric functions.

Keywords : n -additive mappings, Hyers–Ulam–Rassias stability, stability threshold.

MSC (2010) : 39B82; 39B52.

1 Introduction

The control functions of Hyers–Rassias type usually have a threshold of stability, i.e. there is a family of control functions of this type for which stability is flawed. The first and most famous example was given by Gajda [6] for the stability of Cauchy’s equation. Aoki [2], Rassias [7], for $r < 1$, and then Gajda [6], for $r > 1$, showed that if S is a normed vectorial space, B is a Banach space, $\epsilon > 0$, and $f : S \rightarrow B$ is a function such that

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon (\|x\|^r + \|y\|^r), \quad x, y \in S \quad (1)$$

(with the convention that $\|0\|^r = 1$ if $r \leq 0$) then there exists a unique additive function $a : S \rightarrow B$ such that

$$\|f(x) - a(x)\| \leq \epsilon \frac{2}{|2 - 2^r|} \|x\|^r, \quad x \in S.$$

If $r = 1$, the assertion no longer remains valid. Let $f_G : \mathbb{R} \rightarrow \mathbb{R}$,

$$f_G(x) := \sum_{k=0}^{\infty} 2^{-k} \zeta(2^k x), \quad x \in \mathbb{R},$$

where the mapping $\zeta : \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$\zeta(x) := \begin{cases} \frac{\epsilon}{6}, & \text{if } x \in [1, \infty), \\ \frac{\epsilon}{6}x, & \text{if } x \in (-1, 1), \\ -\frac{\epsilon}{6}, & \text{if } x \in (-\infty, -1]. \end{cases}$$

Lemma 1.1. [6] *The function f_G verifies (1), i.e. $|f_G(x+y) - f_G(x) - f_G(y)| \leq \epsilon(|x| + |y|)$, $x, y \in \mathbb{R}$ but if $\delta > 0$ and $m : \mathbb{R} \rightarrow \mathbb{R}$ is an additive mapping, there exists $x_m \in \mathbb{R}$ such that $|f_G(x_m) - m(x_m)| > \delta|x_m|$.*

We agree to say that $r = 1$ is a *stability threshold* for (Hyers -Ulam-Rassias stability of) additive mappings.

The symmetric and n -additive (or multiadditive) functions are important tools in the characterizations of Fréchet polynomials (see, for instance, [1] for a new proof of the famous result of Fréchet [5]).

In this paper we complete the above-mentioned result and we find the threshold of stability for n -additive and symmetric mappings in the particular case of a class of control functions of Hyers-Ulam-Rassias type.

2 Main results

In the following lines we consider that S is an abelian semigroup, B is a Banach space, and n is a positive integer.

For all symmetric function $g : S^n \rightarrow B$, we denote by $D_n g : S^{n+1} \rightarrow B$ the mapping defined by $D_n g(x_1, x_2) := g(x_1, x_2) - g(x_1) - g(x_2)$ for $n = 1$, and for $n > 1$ by

$$D_n g(x_1, \dots, x_{n+1}) := g(x_1, \dots, x_{n-1}, x_n + x_{n+1}) - g(x_1, \dots, x_{n-1}, x_n) - g(x_1, \dots, x_{n-1}, x_{n+1}).$$

We remember that the symmetric function $g : S^n \rightarrow B$ is n -additive if and only if $D_n g = 0$.

If $\varphi : S^{n+1} \rightarrow [0, \infty)$ is a function, $r_n \varphi : S^n \rightarrow [0, \infty)$ denotes the mapping defined by $r_n \varphi(x_1) := \varphi(x_1, x_1)$ if $n = 1$, and for $n > 1$

$$r_n \varphi(x_1, \dots, x_n) := \varphi(2x_1, \dots, 2x_{n-1}, x_n, x_n) + 2\varphi(2x_1, \dots, 2x_{n-2}, x_n, x_{n-1}, x_{n-1}) + \dots + 2^{n-2}\varphi(2x_1, x_n, x_{n-1}, \dots, x_3, x_2, x_2) + 2^{n-1}\varphi(x_n, x_{n-1}, \dots, x_2, x_1, x_1).$$

Let $\varphi : S^{n+1} \rightarrow [0, \infty)$. In [3] we proved that if

$$\sum_{k=0}^{\infty} 2^{-n(k+1)} \varphi_{n+1}(2^k z) < \infty, \quad z \in S^{n+1}, \quad (2)$$

then

$$\tilde{\varphi}^+ := \{\Phi : S^n \rightarrow [0, \infty) \mid \Phi(y) \geq R_n^+ \varphi(y), \text{ and } \lim_{k \rightarrow \infty} 2^{-nk} \Phi(2^k y) = 0, \ y \in S^n\}$$

is a nonempty class, where

$$R_n^+ \varphi(y) := \sum_{k=0}^{\infty} 2^{-n(k+1)} r_n \varphi(2^k y).$$

Also, in [4] we proved that if S is a commutative 2-divisible commutative semigroup and

$$\sum_{k=0}^{\infty} 2^{nk} \varphi_{n+1}(2^{-k-1} z) < \infty, \quad z \in S^{n+1}, \quad (3)$$

then

$$\tilde{\varphi}^- := \{\Phi : S^n \rightarrow [0, \infty) \mid \Phi(y) \geq R_n^- \varphi(y), \text{ and } \lim_{k \rightarrow \infty} 2^{nk} \Phi(2^{-k} y) = 0, \quad y \in S^n\}$$

is a nontrivial class, where

$$R_n^- \varphi(y) := \sum_{k=0}^{\infty} 2^{nk} r_n \varphi(2^{-k-1} y).$$

Using the following elementary lemma, in [3] we have shown that the functions which verify (2) constitute a class of control functions that provide stability for n -additive and symmetric functions.

Lemma 2.1. [3] *Let $(b_k)_{k \in \mathbb{N}}$ be a sequence in B , $(\alpha_k)_{k \in \mathbb{N}}$ be a sequence of positive numbers, and $c > 0$ such that $\beta := \sum_{k=0}^{\infty} c^{-k-1} \alpha_k < \infty$ and $\|b_{k+1} - cb_k\| \leq \alpha_k$, for all $k \in \mathbb{N}$. Then $(c^{-k} b_k)_{k \in \mathbb{N}}$ is a convergent sequence and $\|b - b_0\| \leq \beta$, where $b = \lim_{k \rightarrow \infty} c^{-k} b_k$.*

In the following lines we complete that result using control functions which verifies (3). For convenience, we reproduce from [3] the proof of the result mentioned above also.

Theorem 2.2. *Let $\varphi : S^{n+1} \rightarrow [0, \infty)$ and $g : S^n \rightarrow B$ be a symmetric function satisfying the inequality*

$$\|D_n g(z)\| \leq \varphi(z), \quad z \in S^{n+1}. \quad (4)$$

1. If φ verifies (2) then there exists a unique symmetric n -additive function $a : S^n \rightarrow B$ such that

$$\|g(y) - a(y)\| \leq \Phi(y), \quad y \in S^n \quad (5)$$

for all $\Phi \in \tilde{\varphi}^+$. The symmetric n -additive function a is defined by $a(y) := \lim_{k \rightarrow \infty} 2^{-nk} g(2^k y)$, $y \in S^n$.

2. If S is a 2-divisible abelian semigroup and φ verifies (3), then there exists a unique symmetric n -additive function $a : S^n \rightarrow B$ which satisfies (5) for all $\Phi \in \tilde{\varphi}^-$. The symmetric n -additive function a is defined by $a(y) := \lim_{k \rightarrow \infty} 2^{nk} g(2^{-k} y)$, $y \in S^n$.

Proof. Let $y = (x_1, \dots, x_n) \in S^n$. Putting $z = (x_1, \dots, x_{n-1}, x_n, x_n)$ in (5), we get

$$\begin{aligned} \|D_n g(x_1, \dots, x_{n-1}, x_n, x_n)\| &= \|g(x_1, \dots, x_{n-1}, 2x_n) - 2g(x_1, \dots, x_n)\| \leq \\ &\leq \varphi(x_1, \dots, x_{n-1}, x_n, x_n). \end{aligned}$$

But g is a symmetric function; therefore,

$$\begin{aligned} \|g(2y) - 2^n g(y)\| &= \| [g(2x_1, 2x_2, \dots, 2x_{n-1}, 2x_n) - 2g(2x_1, 2x_2, \dots, 2x_{n-1}, x_n)] + \\ &+ 2[g(2x_1, 2x_2, \dots, 2x_{n-2}, x_n, 2x_{n-1}) - 2g(2x_1, 2x_2, \dots, 2x_{n-2}, x_n, x_{n-1})] + \dots \\ &+ 2^{n-1} [g(x_n, x_{n-1}, \dots, x_2, 2x_1) - 2g(x_n, x_{n-1}, \dots, x_2, x_1)] \| \leq \\ &\|D_n g(2x_1, \dots, 2x_{n-1}, x_n, x_n)\| + 2\|D_n g(2x_1, \dots, 2x_{n-2}, x_n, x_{n-1}, x_{n-1})\| + \dots \\ &+ 2^{n-1} \|D_n g(x_n, x_{n-1}, \dots, x_2, x_1, x_1)\| \leq r_n \varphi(x_1, \dots, x_n), \end{aligned}$$

hence,

$$\|g(2y) - 2^n g(y)\| \leq r_n \varphi(y), \quad y \in S^n. \quad (6)$$

1. Suppose that φ verifies (2) and that $\Phi \in \tilde{\varphi}^+$. Replacing y by $2^k y$ ($k \in \mathbb{N}$) in (6), we get

$$\|g(2^{k+1} y) - 2^n g(2^k y)\| \leq r_n \varphi(2^k y), \quad y \in S^n.$$

Applying Lemma 2.1 (for $b_k = g(2^k y)$, $c = 2^n$, $\alpha_k = r_n \varphi(2^k y)$ and $\beta = R_n^+ \varphi(y) \leq \Phi(y)$), it follows that $(2^{-nk} g(2^k y))_{k \in \mathbb{N}}$ is a convergent sequence in B , and its limit, $a(y) := \lim_{k \rightarrow \infty} 2^{-nk} g(2^k y)$, satisfies (5). Since g is symmetric, it follows that a is a symmetric function, too. From (4) and (2) it follows that

$$\lim_{k \rightarrow \infty} 2^{-nk} \|D_n g(2^k z)\| \leq \lim_{k \rightarrow \infty} 2^{-nk} \varphi(2^k z) = 0,$$

whence $D_n a(z) = 0$, $z \in S^{n+1}$, i.e. a is a symmetric and n -additive mapping which satisfies (5). If $a' : S^n \rightarrow B$ is an n -additive mapping and

$$\|g(y) - a'(y)\| \leq \Phi(y), \quad y \in S^n,$$

since $a'(2^k y) = 2^{nk} a'(y)$, and $\Phi \in \tilde{\varphi}^+$, we have

$$\lim_{k \rightarrow \infty} \|2^{-nk} g(2^k y) - a'(y)\| \leq \lim_{k \rightarrow \infty} 2^{-nk} \in \Phi(2^k y) = 0, \quad y \in \mathcal{S}^n,$$

whence $a' = a$; therefore a is the unique symmetric and n -additive mapping which satisfies (5).

2. Suppose now that φ verifies (3) and that $\Phi \in \tilde{\varphi}^-$. Replacing y by $2^{-k-1}y$ ($k \in \mathbb{N}$) in (6), we get

$$\|g(2^{-k-1}y) - 2^{-n}g(2^{-k}y)\| \leq 2^{-n}r_n\varphi(2^{-k-1}y), \quad y \in \mathcal{S}^n.$$

Using again Lemma 2.1 (for $b_k = g(2^{-k}y)$, $c = 2^{-n}$, $\alpha_k = 2^{-n}r_n\varphi(2^{-k-1}y)$ and $\beta = R_n^-\varphi(y) \leq \Phi(y)$), it follows, as in the first case, that $a(y) := \lim_{k \rightarrow \infty} 2^{nk}g(2^{-k}y)$ defines the unique symmetric n -additive mapping which satisfies (5). $\square$

The next consequence is a stability result in the Aoki-Rassias sense.

Corollary 2.3. Let S be a normed space, $\epsilon > 0$ and $r \neq 1$. Suppose that $g : S^n \rightarrow B$ is a symmetric function such that

$$\|D_n g(x_1, \dots, x_{n+1})\| \leq \epsilon \|x_1\|^r \cdots \|x_{n-1}\|^r (\|x_n\|^r + \|x_{n+1}\|^r) \quad (7)$$

for all $x_1, \dots, x_{n+1} \in S$. Then there exists a unique n -additive mapping $a : G^n \rightarrow B$ for which

$$\|g(x_1, \dots, x_n) - a(x_1, \dots, x_n)\| \leq \epsilon \frac{2^{(n-1)(r-1)+1}}{|2^r - 2|} \|x_1\|^r \cdots \|x_n\|^r, \quad (8)$$

for all $x_1, \dots, x_n \in S \setminus \{0\}$. If $r < 1$ then $a(y) := \lim_{k \rightarrow \infty} 2^{-nk}g(2^k y)$, and if $r \in (1, \infty)$, then $a(y) := \lim_{k \rightarrow \infty} 2^{nk}g(2^{-k}y)$, $y \in S^n$.

Proof. Let

$$\varphi(x_1, \dots, x_{n+1}) := \epsilon \|x_1\|^r \cdots \|x_{n-1}\|^r (\|x_n\|^r + \|x_{n+1}\|^r).$$

Then

$$r_n\varphi(x_1, \dots, x_n) = \frac{2^{(n-1)(r-1)+1}(2^{nr} - 2^n)}{2^r - 2}.$$

1. Let $r < 1$. Then φ verifies (2). But, for $x_1, \dots, x_n \neq 0$, we have

$$R_n^+\varphi(x_1, \dots, x_n) = \frac{r_n\varphi(x_1, \dots, x_n)}{2^{nr} - 2^n} = \frac{2^{(n-1)(r-1)+1}}{2^r - 2} \|x_1\|^r \cdots \|x_n\|^r.$$

We apply Theorem 2.2 for $\Phi = R_n^+ \varphi$ and we obtain (8) for $a(y) = \lim_{k \rightarrow \infty} 2^{-nk} g(2^k y)$.

2. Let $r > 1$. Then φ verifies (3) and

$$R_n^- \varphi(x_1, \dots, x_n) = \frac{r_n \varphi(x_1, \dots, x_n)}{2^n - 2^{nr}} = \frac{2^{(n-1)(r-1)+1}}{2 - 2^r} \|x_1\|^r \cdots \|x_n\|^r.$$

We apply Theorem 2.2 for $\Phi = R_n^- \varphi$ and we obtain (8) for $a(y) := \lim_{k \rightarrow \infty} 2^{nk} g(2^{-k} y)$, $y \in S^n$. $\square$

Stability threshold is $r = 1$

Let $S = B = \mathbb{R}$ and $n \geq 2$. Let $\epsilon > 0$. The stability problem for n -additive and symmetric mappings in the case $r = 1$ is: there exists a positive constant δ such that if $g : \mathbb{R}^n \rightarrow \mathbb{R}$ is a symmetric function and

$$|D_n g(x_1, \dots, x_n, x_{n+1})| \leq \epsilon |x_1| \cdots |x_{n-1}| (|x_n| + |x_{n+1}|), \quad x_1, \dots, x_{n+1} \in \mathbb{R}, \quad (9)$$

there exists a unique symmetric and n -additive mapping $a : \mathbb{R}^n \rightarrow \mathbb{R}$ for which

$$|g(x_1, \dots, x_n) - a(x_1, \dots, x_n)| \leq \delta |x_1| \cdots |x_n|, \quad x_1, \dots, x_n \in \mathbb{R}. \quad (10)$$

We give two examples. The first: a function g which verifies (9), but for which there exist an infinity of symmetric and n -additive mappings satisfying (10). The second: a function g which verifies (9), but for which there does not exist a symmetric and n -additive mapping a satisfying (10).

1. The symmetric function g defined by $g(x_1, \dots, x_n) := \frac{\epsilon}{2} |x_1| \cdots |x_n|$ verifies (9), and, for all $\alpha \in [-\delta + \frac{\epsilon}{2}, \delta + \frac{\epsilon}{2}]$ $a(x_1, \dots, x_n) := \alpha x_1 \cdots x_n$ defines a symmetric n -additive mapping which satisfies (10).

2. From Lemma 1.1, it follows that the function $f_G : \mathbb{R} \rightarrow \mathbb{R}$ verifies

$$|f_G(x+y) - f_G(x) - f_G(y)| \leq \epsilon (|x| + |y|), \quad x, y \in \mathbb{R} \quad (11)$$

and, for all additive mapping $m : \mathbb{R} \rightarrow \mathbb{R}$, there exists $x_m \in \mathbb{R}$ for which

$$|f_G(x_m) - a(x_m)| > \delta |x_m|. \quad (12)$$

Let and $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be the symmetric function defined by

$$g(x_1, \dots, x_n) := f_G(x_1) x_2 \cdots x_n + x_1 f_G(x_2) x_3 \cdots x_n + \cdots + x_1 \cdots x_{n-1} f_G(x_n).$$

From (11) it follows that g satisfies (9). Suppose that $a : \mathbb{R}^n \rightarrow \mathbb{R}$ is a symmetric and n -additive mapping which verifies (10). Then, for $x_1 = x_2 = \dots = x_{n-1} = 1$, and $x_n = x$, we have from (10):

$$|f_G(x) - [a(1, \dots, 1, x) - (n-1)f_G(1)x]| \leq \delta|x|, \text{ for all } x \in \mathbb{R}. \quad (13)$$

But $m(x) := a(1, \dots, 1, x) - (n-1)f_G(1)x$ defines an additive mapping and therefore (13) contradicts (12). $\square$

References

- [1] J. M. Almira, A. J.- López-Moreno, *On solutions of the Fréchet functional equation*, J. Math. Anal. Appl., **332:2**, (2007), doi:10.1016/j.jmaa.2006.11.002, 1119-1133
- [2] T. Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan **2** (1950), 84-88.
- [3] D. M. Dăianu, *Recursive procedure in the stability of Fréchet polynomials*, Adv. Difference Equ. **2014:16** (2014), doi:10.1186/1687-1847-2014-16.
- [4] D. M. Dăianu, *A stability criterion for Fréchet's first polynomial equation*, Aequationes Math. (2014), DOI: 10.1007/s00010-014-0257-7.
- [5] M. Fréchet, *Une définition fonctionnelle des polynômes*, Nouv. Ann. Math. **9** (1909), 145-182.
- [6] Z. Gajda, *On stability of additive mappings*, Int. J. Math. Sci. **14** (1991), 431-434.
- [7] Th.M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), 297-300.

Dan M. Dăianu
 Department of Mathematics,
 "Politehnica" University of Timișoara,
 Victoriei Square, No.2,
 300006 Timișoara, Romania,
 dan.daianu@upt.ro